


How effective are created or restored freshwater wetlands for nitrogen and phosphorus removal?

A systematic review protocol

Magnus Land
Wilhelm Granéli
Anders Grimvall
Carl Christian Hoffmann
William Mitsch
Karin Tonderski
Jos Verhoeven



Mistra EviEM SR2 Protocol

Publication date: 28 August 2013

This protocol follows the CEE *Guidelines for systematic reviews in environmental management*, issued by the [Collaboration for Environmental Evidence](#).

It has also been published in the open-access journal [Environmental Evidence](#).

Please cite as: Land *et al.*: How effective are created or restored freshwater wetlands for nitrogen and phosphorus removal? A systematic review protocol. *Environmental Evidence* 2013, **2**:16.

How effective are created or restored freshwater wetlands for nitrogen and phosphorus removal?

A systematic review protocol

Magnus Land^{1*}

* Corresponding author

Email: magnus.land@eviem.se

Wilhelm Granéli²

Email: wilhelm.graneli@biol.lu.se

Anders Grimvall³

Email: anders.grimvall@havsmiljoinstitutet.se

Carl Christian Hoffmann⁴

Email: cch@dmu.dk

William Mitsch⁵

Email: wmitsch@fgcu.edu

Karin Tonderski⁶

Email: karsu@ifm.liu.se

Jos Verhoeven⁷

Email: j.t.a.verhoeven@uu.nl

1 Mistra EviEM, The Royal Swedish Academy of Sciences, Box 50005, SE–104 05 Stockholm, Sweden

2 Lund University, Dept Biol, SE–22362 Lund, Sweden

3 Swedish Institute for the Marine Environment, Box 260, S–405 30 Gothenburg, Sweden

4 Aarhus University, Dept Biosci, DK–8600 Silkeborg, Denmark

5 110 Kapnick Center, Florida Gulf Coast University, 4940 Bayshore Drive, Naples, Florida 34112, USA

6 Linköping University, Dept Phys, Chem and Biol, SE–581 83 Linköping, Sweden

7 Utrecht University, Dept Biol, NL–3584 CA Utrecht, Netherlands

Abstract

Background

Eutrophication of aquatic environments is a major environmental problem in large parts of the world. In Europe, EU legislation (the Water Framework Directive and the Marine Strategy Framework Directive), international conventions (OSPAR, HELCOM) and national environmental objectives emphasize the need to reduce the input of plant nutrients to freshwater and marine environments. A widely used method to achieve this is to let water pass through a constructed or restored wetland (CW). However, the large variation in measured nutrient removal rates in such wetlands calls for a systematic review. The objective of this review is to quantify nitrogen and phosphorus removal rates in constructed or restored wetlands and relate them to wetland characteristics, loading characteristics, and climate factors. Wetlands are created to treat water from a number of different sources. Sources that will be considered in this review include agricultural runoff and urban storm water run-off, as well as aquaculture wastewater and outlets from domestic wastewater treatment plants, with particular attention to the situation in Sweden. Although the performance of wetlands in temperate and boreal regions is most relevant to the Swedish stakeholders a wider range of climatic conditions will be considered in order to make a thorough evaluation of climatic factors.

Methods

Searches for primary studies will be performed in electronic databases as well as on the internet. One author will perform the screening of all retrieved articles at the title and abstract level. To check that the screening is consistent and complies with the agreed inclusion/exclusion criteria, subsets of 100 articles will be screened by the other authors. When screening at full-text level the articles will be evenly distributed among the authors. Kappa tests will be used to evaluate screening consistency. Data synthesis will be based on meta-regression. The nutrient removal rates will be taken as response variables and the effect modifiers will be used as explanatory variables. More specifically, the meta-regression will be performed using generalized additive models that can handle nonlinear relationships and major interaction effects. Furthermore, subgroup analyses will be undertaken to elucidate statistical relationships that are specific to particular types of wetlands.

Keywords

Nitrogen, Phosphorus, Nutrient, Retention rate, Removal efficiency, Wetland creation, Restored wetland, Constructed wetland, Pond, Eutrophication

Background

Man-made eutrophication of aquatic environments is a major environmental problem in large parts of the world [1]. This is also the case in Europe, where EU legislation (the Water Framework Directive and the Marine Strategy Framework Directive), international conventions (OSPAR, HELCOM) and national environmental objectives emphasize the need to reduce the input of plant nutrients to freshwater and marine environments. Furthermore, it is generally accepted that some of the reduction measures target agricultural runoff and other diffuse emissions of nitrogen and phosphorus.

One method to reduce the input of plant nutrients to freshwater and coastal waters is to let water pass through some kind of constructed or restored wetland. Major types of constructed wetlands include Free Surface Flow Constructed Wetlands, Subsurface Horizontal Flow Constructed Wetlands, and Subsurface Vertical Flow Constructed Wetlands, whereas restored wetlands refer to wetlands where interventions have been made to recreate previously drained or by other means altered natural wetlands. The physical and biogeochemical processes involved in the removal of nitrogen and phosphorus in such wetlands are relatively well known. However, the large variation in efficiency observed in different studies calls for a systematic review of removal rates and how they are influenced by the wetland characteristics, loading differences, and environmental factors. For non-point source runoff, Mitsch *et al.* [2] found an average nitrogen reduction of $39 \text{ g m}^{-2} \text{ yr}^{-1}$ in Ohio wetlands and $46 \text{ g m}^{-2} \text{ yr}^{-1}$ in a large wetland in Louisiana, and attributed the difference to the warmer climate in the latter region. Carleton *et al.* [3] analysed time series from 49 wetlands treating storm water runoff and found that nitrogen removal (% of load) increased in a non-linear way with an increasing ratio wetland/catchment area. In another data set from 65 constructed wetlands, the nitrate removal varied from around 1 to $> 1000 \text{ g m}^{-2} \text{ yr}^{-1}$, and the variation was best explained by a first order equation including inflow concentration, load, temperature and hydraulic efficiency as explaining variables [4].

Removal rates for wetlands treating wastewater are generally higher. Vymazal [5] reviewed nitrogen retention in constructed wetlands (Free-floating Plants (FFP), Free Water Surface systems (FWS), horizontal sub-surface flow systems (HSSF) and vertical sub-surface flow systems (VSSF)) and concluded that removal of total nitrogen varied between 40 and 50% with removed load ranging between 250 and $630 \text{ g N m}^{-2} \text{ yr}^{-1}$ depending on CW type and inflow loading.

Vymazal and Kroepfelova [6] described the removal of nitrogen in horizontal sub-surface flow constructed wetlands (HF CWs) for various types of wastewater. The survey included more than 900 annual means from more than 300 systems in 36 countries that were categorized into municipal, industrial, agricultural, and landfill leachate. While the median inflow load of total nitrogen was $574 \text{ g m}^{-2} \text{ yr}^{-1}$, the median removal rate was $225 \text{ g m}^{-2} \text{ yr}^{-1}$.

Regarding phosphorus, Carleton *et al.* [3] concluded that removal of total phosphorus was more a function of mean detention time than of mean hydraulic loading rate in wetlands receiving storm water runoff. Hoffmann *et al.* [7] found phosphorus retention varying from a loss of $1 \text{ g m}^{-2} \text{ yr}^{-1}$ to a retention rate of $0.3 \text{ g m}^{-2} \text{ yr}^{-1}$ in wetlands and floodplains. Losses were often in the form of dissolved phosphate. In contrast, Braskerud *et al.* [8] reported retention of $27 - 156 \text{ g m}^{-2} \text{ yr}^{-1}$ phosphorus in small constructed wetlands in Norway, and could attribute the high retention to settling of particulate phosphorus. Without management activities phosphorus removal may decrease over time, as was indicated by Mitsch *et al.* [9] who showed that TP retention dropped from an initial 60% to 10% 15 years after wetland construction.

In Sweden, the main focus when wetlands initially started to be constructed and restored was on nitrogen retention. Nitrogen was usually assumed to limit primary production in marine ecosystems [10,11], and also in the brackish-water Baltic Sea [12] which catchment includes most of Sweden. However, this is a disputed assumption, and some scientists have the opinion that phosphorus is ultimately limiting production in the Baltic Sea [13-15]. In freshwater bodies eutrophication is usually thought to be controlled by phosphorus inputs only [14], although this is still somewhat controversial and some scientists argue that also nitrogen inputs to lakes have to be reduced as well [13]. Thus, to single out one nutrient or the other as limiting in the marine and freshwater system, respectively, may simplify reality too far. In many systems both nitrogen and phosphorus are limiting depending on time of year and location [16], and therefore, whether the major concern is the marine environment or freshwater ecosystems, quantification of both nitrogen and phosphorus retention is very relevant.

In Sweden, removal of nitrogen and phosphorus in CWs is of particular interest because such measures may play an important role to achieve national environmental quality objectives and to reach the commitments made in the Baltic Sea Action Plan (BSAP). A large number of wetlands have been created; e.g. during the period 2000–2012 more than 1400 wetlands were supported by various governmental funds [17]. The performance of some of them has been evaluated by Svensson *et al.* [18] who reported that the nitrogen retention rates varied an order of magnitude depending on which subsidy program that had funded the wetland, and that they ranged between 5 and 54 g m⁻² y⁻¹. The variations were mainly attributed to differences in nitrogen loading rates.

The consortium Swedish Environmental Emissions Data (SMED) has used a catchment model to evaluate the effect of >4000 ha created/restored wetlands on the transport of nitrogen and phosphorus to the Baltic Sea from Sweden south of River Dalälven [19]. In the report, the average nitrogen and phosphorus removal rates were calculated to 3.4 and 0.3 g m⁻² y⁻¹, respectively, but with a considerable uncertainty.

Weisner and Thiere [20] used essentially the same calculation methods as Svensson *et al.* [18] to compare the nitrogen and phosphorus retention in wetlands constructed before and after 2003, respectively. For all wetlands constructed in the period 2003–2008 (n = 50) the result was 5.9-10 g m⁻² y⁻¹ for nitrogen and 0.17 – 0.53 g m⁻² y⁻¹ for phosphorus. In contrast, the removal rates were 59–65 g m⁻² y⁻¹ for nitrogen and 0.68 – 1.5 g m⁻² y⁻¹ for phosphorus in wetlands located in the county of Skåne (n = 6), where the losses of nutrients from farmland are high.

The mentioned Swedish studies are largely based on model calculations on various scales, and although the results are not directly comparable, it appears that the calculated rates of nitrogen and phosphorus retention in constructed or restored wetlands span over a wide range. It is thus not quite clear to what degree constructed and restored wetlands will contribute to fulfilling the Swedish commitments in the BSAP, or how important they are for mitigating eutrophication in freshwater bodies. In this context, The Swedish Board of Agriculture, which administers a major part of the governmental funds for wetland construction and restoration, and the Swedish Agency for Marine and Water Management, which is responsible for the environmental goal “zero eutrophication”, were interested to obtain better information on measured retention rates in individual wetlands. They were also interested in the factors controlling the difference in performance between different wetlands. The idea is that getting a coherent picture of how different wetlands function in a variety of conditions should make it easier to plan more effective water pollution control. Another important stakeholder in this systematic review is the wastewater treatment industry, which in the future most likely will be expected to comply to even heavier regulations on nutrient emissions than today.

Objectives of the review

The objective of this review is to quantify observed retention rates of nutrients in constructed or restored wetlands, and also to examine the distribution of these rates and quantify the variation between different studies. The primary question this review seeks an answer to is “How effective are created or restored freshwater wetlands for nitrogen and phosphorus removal?”

Secondary questions are related to how various effect modifiers, such as environmental conditions and wetland characteristics, influence the nutrient removal rates. For that reason this review covers a fairly wide range of climatic conditions, although the performance of wetlands in temperate and boreal regions is most relevant to the stakeholders in Sweden. The review will not engage in detailed investigations of various removal processes and mechanisms but rather treat each wetland as a “black box”. Such an approach may of course introduce some uncertainties. However, when assessing study quality, studies presenting complete nutrient budgets where removal by individual processes have been quantified will be rated higher than studies merely providing inlet/outlet data. A balanced budget may indicate that the numbers are reasonable accurate and/or that no major source or sink have been overlooked.

The structure of the primary question is further discussed in the section Study inclusion criteria where information on relevant subjects, interventions, comparator and outcomes is given in more detail.

Methods

Searches

Searches in electronic databases will be made using the search terms displayed in Table 1. Using the Boolean operators indicated this translates into the search string below, where * is a wildcard that can be any number of characters. Different wildcards may apply in different databases and modifications will be made accordingly.

Table 1. Suggested search strategy

	Term 1 (intervention)	AND	Term 2 (intervention)	AND	Term 3 (subject)	AND	Term 4 (outcome)
	wetland*		construct*		nitrogen		retention
OR	pond	OR	create*	OR	phosph*	OR	trap*
OR	mire*	OR	restor*	OR	nitrate	OR	denitrification
OR	marsh	OR	Man*made	OR	TKN	OR	uptake
OR	fen	OR	flooding	OR	ammoni*	OR	sedimentation

OR	wet meadow	OR	inundation	OR	remov*
OR	riparian	OR	artificial	OR	settling
OR	flood plain			OR	accretion
OR	reed			OR	precipitation
				OR	*sorption

Search terms are connected with AND, and words within each term is connected by OR. Truncation or wildcard is denoted by *.

(Wetland* OR pond OR mire* OR marsh OR fen OR “wet meadow” OR riparian OR “flood plain” OR reed) AND (construct* OR creat* OR restor* OR man*made OR flooding OR inundation OR artificial) AND (nitrogen OR phosph* OR nitrate OR TKN OR ammoni*) AND (retention OR trap* OR denitrification OR uptake OR sedimentation OR remov* OR settling OR accretion OR precipitation OR *sorption).

No particular time, document type or language constraints will be applied. Electronic databases that will be used for searching are listed in Table 2. The table also indicates the fields that will be searched and number of hits obtained in preliminary searches.

Table 2. Electronic databases used for searching

Database	Searched field	No of hits	Date
ISI Web of Science	topic	3249	2013-02-26
Georef and Geobase	subject/title/abstract	2249	2013-02-26
Scopus	title, abstract and keywords	2842	2013-02-26
Agricola	keyword anywhere	524	2013-02-26
ASFA	All fields (no full text)	1933 ^{a)}	2013-02-27
Academic search	title/abstract/subject/keyword	1131	2013-02-27
Biological Abstracts	keywords	401	2013-02-27
Wiley Online Library	abstract or title or keywords	174	2013-02-27
Directory of Open Access Journals	all fields	84 ^{a)}	2013-03-05
ScienceDirect	abstract or title or keywords	876	2013-07-02

^{a)} After removal of duplicates.

The search engine in Directory of Open Access Journals (DOAJ) does not allow complex search strings. Therefore the search strategy in that data base has to be simplified. The word “wetland” will be

combined with each of the subject words (term 3 in Table 1) using the Boolean operator AND. Since no truncations are allowed in the DOAJ search engine, separate searches will be done using the words phosphorus and phosphate as well as ammonium and ammonia.

The searches are expected to retrieve a number of review articles not providing any original data. Even though such articles will not be used for data extraction, their bibliographies will be used to estimate the comprehensiveness of the searches.

In addition to data in the scientific literature it is anticipated that data will be found also in the grey literature. Such data will be searched for on the internet. When searching the internet the search terms will be in English as well as in Swedish, Danish and Dutch. Searches on the internet will be performed using Google Scholar where the simplified search strings shown below will be applied (the search string used for the electronic databases is too long for Google scholar).

English: (wetland OR pond OR marsh) AND (constructed OR created OR restored OR artificial) AND (nitrogen OR phosphorus) AND (retention OR removal).

Swedish: (våtmark OR damm) AND (anlagd OR skapad OR restaurerad OR artificiell) AND (kväve OR fosfor) AND (retention OR rening OR avskiljning).

Danish: (vådområde OR lavvandet sø OR minivådområde) AND (konstrueret OR reetableret OR genskabt OR kunstig) AND (kvælstof OR nitrogen OR fosfor OR fosfat) AND (tilbageholdelse OR retention OR fjernelse OR rensning).

Dutch: (moeras OR rietveld OR sloot) AND (kunstmatig OR aangelegd OR hersteld OR helofytenfilter) AND (stikstof OR fosfaat OR fosfor) AND (verwijdering OR zuivering).

In addition, websites of relevant specialist organisations (listed below) will also be searched, using the same search strings (in relevant languages) as for Google scholar. In general the first 100 hits will be examined for internet searches.

- Swedish Environmental Protection Agency (SEPA)
- Swedish Board of Agriculture
- The Swedish Agency for Marine and Water Management
- Swedish directory of Master thesis (DiVA)
- South Florida Water Management District
- U.S. Environmental Protection Agency (EPA)
- North American Data Base (NADB)
- U.S. Department of Agriculture (USDA)
- Foundation for Applied Water Research (STOWA)
- Ekologgruppen i Landskrona AB
- Norwegian Institute for Agricultural and Environmental Research (Bioforsk)
- Danish Centre for Environment and Energy (DCE)
- European Environment Agency (EEA)
- Wetland Solutions Inc.
- Wetlands International
- Finnish Environment Institute (SYKE)
- Federal Environment Agency (UmweltBundesamt, Germany)
- Stichting Toegepast Waterbeheer (STOWA, The Netherlands)

Study inclusion/exclusion criteria

Wetlands are created to treat water from a number of different sources. Sources that will be considered in this review include agricultural runoff and urban storm water run-off, as well as outlets from domestic wastewater treatment plants. In the statistical analysis of data they will however be treated separately. Untreated wastewater will not be considered since in most countries in Europe it is not allowed to discharge such water into the environment. Moreover, industrial and agricultural wastewater may vary considerably in composition and will therefore be excluded as well.

To be included in the systematic review, each article must pass each of the following criteria. The criteria have been developed in collaboration with stakeholders. In case it cannot be decided on the title and abstract level the article will pass to the full text level.

- *Relevant subject:* Secondary and tertiary treated domestic wastewater, urban storm water, stream/river water, freshwater aquaculture effluents, and agricultural runoff.
- *Types of intervention:* Creation or restoration of wetlands. Created wetlands include both horizontal and vertical subsurface flow systems, as well as free surface flow systems. In addition, there are more or less natural wetlands including riparian wetlands that have been restored in order to enhance either nitrogen and phosphorus retention or biodiversity. Restoration refers to recovery of ecological and hydrological processes as well as geomorphology in areas where natural wetlands previously have been drained or by other means altered. To be included the created or restored wetlands must host some type of vegetation.
- *Types of comparator:* No intervention (inlet conditions can serve as control)
- *Types of outcome:* Removal of total nitrogen and total phosphorus from the water body per unit wetland area and year. Removal efficiency (% of load).
- *Types of study:* The most common way to evaluate the overall retention rate in a wetland is to compare the nitrogen/phosphorus loads in the inlet water and in the outlet water, respectively. Quite often the retention in wetlands is evaluated in experiments where effect modifiers such as loading rate or vegetation type are varied. This is a version of a Control-Impact (CI) study where inlet conditions serve as control. In rare cases nutrient loads in a river or stream have been recorded both before and after the establishment of a wetland, which corresponds to a typical Before-After (BA) study. Both types of studies will be included.

The removal rates and efficiencies may show large seasonal variations. Therefore, to be included in the review, it is a prerequisite that the wetland is established in field conditions and exposed to the ambient climate. This means that laboratory studies will be excluded and that each study must cover at least one complete annual cycle. Also, in order to reflect realistic conditions the wetland must be of a reasonable size. While microcosm studies will be excluded, mesocosm studies will be included since they provide valuable information on the variability of the outcomes based on true replicates, and mesocosms are judged to be of sufficient size.

The review will focus on boreal and temperate regions, but for comparison sub-tropical regions will also be included. In the Köppen-Geiger climate classification system [21] this corresponds roughly to group D (snow climates), group C (warm temperate climates) and parts of group A (Equatorial climates with one dry season, i.e. As and Aw).

Wetlands may be constructed or restored for other purposes than nutrient retention. Although some wetlands serve multiple purposes [22], sometimes the main purpose is to promote biodiversity or reduce flood risks. In this review wetlands will be considered regardless of the main purpose of the

wetland, i.e. inclusion and exclusion will not be based on the reasons for constructing or restoring the wetlands. When data are extracted, however, the main purpose will be recorded.

At the title and abstract level one author will perform the screening of all retrieved articles. To check that the screening is consistent and complies with the agreed inclusion/exclusion criteria, a subset of 100 articles will also be screened by another author. A second subset of 100 articles will be screened by two other authors and a third subset of articles will be screened by another pair of authors. In this way 300 articles will be double-screened. It will also be possible to check the consistency between the main screener and the other authors as well as between the other authors within each screening pair. To evaluate the consistency Kappa tests will be used.

When screening at the full-text level the articles will be evenly distributed among the authors. However, before screening at full scale, a subset of approximately 100 of the articles will be double-screened. Again, Kappa tests will be used to test the consistency between the authors.

If any Kappa test shows unacceptable discrepancies ($\kappa < 0.6$) the inclusion/exclusion criteria will be revisited by all authors and formulated in a less ambiguous way.

Potential effect modifiers and reasons for heterogeneity

The nutrient retention may vary considerably between different studies. The anticipated large variation is easy to understand in the light of the fact that the retention rate is a result of several independent parameters. Nitrogen removal takes place through 1) sedimentation, 2) plant uptake, and 3) denitrification and volatilization. Processes involved in phosphorus removal are 1) sedimentation, 2) plant uptake 3) sorption and 4) precipitation/co-precipitation. The success of each of these mechanisms may depend on factors such as:

Loading characteristics

- Hydraulic loading rate
- Concentration and speciation of nitrogen and phosphorus at the inlet

Wetland characteristics

- Type of wetland
- Size and shape (area, depth, length)
- Hydroperiod
- Age
- Sediment/soil type
- Oxygen concentration and redox potential
- Vegetation type and coverage
- Fauna
- Management methods and frequency

Climate characteristics

- Mean temperature
- Ice coverage

Study quality assessment

Studies that still are included after the full text screening will be subject to a quality assessment. During this process the studies will be assigned to either of three quality categories: 1) Does not meet the quality criteria, 2) Acceptable, and 3) High standard. Studies in category 1 will not qualify for meta-analysis. Data may however be extracted and compared to studies that are assigned to higher quality categories. In the same way the results from category 2 studies will be compared to the results from category 3 studies. The intention is to examine if there are any significant differences and, if so, low-quality studies may provide misleading information. Other than that there will be no further weighting of the studies.

The quality assessment will focus on the following aspects:

- A study length of 12 months is acceptable. Longer study periods will be regarded as higher quality. In an earlier review on natural wetlands [23], it was shown that studies conducted over a period of a year or more, or that involved frequent sampling during high flow events, were more likely to indicate that the wetland increased the nutrient loadings.
- Sampling frequency must be sufficient. A guide-line value of 12 sampling occasions/year is acceptable. Higher sampling frequencies, as well as flow proportional sampling, will render a higher quality rating. Lower sampling frequencies are acceptable in e.g. sedimentation studies where the samples better integrate removal processes over time.
- Hydrological data adequately covering the major water flow pathways should have been used in the calculation of nutrient loads.
- The quality of studies presenting complete nutrient budgets where removal by individual processes are quantified will be regarded as higher than the quality of studies merely providing inlet/outlet data.
- Studies with replicate wetlands will be given a higher quality rating than studies without replicates.
- Studies should account for at least one of the potential effect modifiers related to loading characteristics, wetland characteristics and climate characteristics, respectively (mentioned in previous section).

Data extraction strategy

The basic outcome of the reviewed wetland application consists of a change in nitrogen or phosphorus concentration in the water. However, the outcome this review is seeking to evaluate is the removal rate and removal efficiency; typically the results are reported quantitatively as $\text{g m}^{-2} \text{y}^{-1}$ and as % of load, respectively. Results of studies reported in other units will be recalculated if possible.

In order to assess the quality of the studies and to be able to evaluate the importance of various effect modifiers, other data than the retention rates will be recorded as well (see Table 3). When relating retention rates to the wetland area it is important to make a distinction between the actual area and the effective area, and to account for possible channel effects. In case no annual average retention rates have been calculated, data will be extracted for each sampling occasion and then an annual average will be calculated if possible. To make the data extraction as consistent as possible the data will be entered into a predesigned Excel spreadsheet.

Table 3. Parameters and data that will be recorded if provided

Loading characteristics	Wetland characteristics	Climate characteristics	Study design parameters	Results
Type of water	Type of wetland	Annual mean air temperature	Study length	Annual average removal rates ($\text{g m}^{-2} \text{y}^{-1}$)
Concentrations and speciation of N and P at inlet	Length, Width, Depth	Annual mean water temperature	Sampling frequency	Annual average removal efficiency (% of load)
Annual average inflow loadings	Area	Ice coverage	Flow proportional Sampling (Y/N)	Number of observations (years)
Hydraulic load (m/d)	Hydroperiod (d)	GPS coordinates		Standard deviation
Timing of peak discharge	Dominant plant species and coverage	Climate type ¹⁾		
Residence time (d)	Sediment/soil type			
pH	Land use history and wetland age			
	O ₂ concentration and redox potential			
	Fauna			
	Management methods			
	Main purpose of wetland			
	Other characteristics ²⁾			

¹⁾ According to the Köppen-Geiger climate classification [21]. This data will in most cases not be provided by the primary studies but will be extracted elsewhere.

²⁾ May include, e.g., specific substrates, soil amendments, configuration of multiple wetlands, or other means of enhancing the outcome.

All studies do not provide information on all parameters shown in Table 3. For instance, we do not expect that data on the fauna in the wetlands is reported very frequently. It is however quite possible that the nutrient cycling and retention are influenced by, e.g., benthic organisms through bioturbation [24] or by birds [25,26], and if such data is provided it will be extracted.

Data synthesis and presentation

The data synthesis will be based on meta-regression. The removal rates of nitrogen and phosphorus will be taken as response variables and the effect modifiers listed above will be used as explanatory variables. More specifically, the meta-regression will be performed using generalized additive models that can handle non-linear relationships and major interaction effects. Because the number of potentially important effect modifiers is fairly large, cross-validation will be employed to identify a subset of major effect modifiers. Furthermore, subgroup analyses will be undertaken to elucidate statistical relationships that are specific to particular types of wetlands. Tree analyses will be performed to create an overview of heterogeneous data and nonlinear relationships.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

This systematic review protocol is based on a draft written by ML. All authors discussed the draft and suggested improvements at a meeting in December 2012. The draft was then finalized by ML with significant contributions from AG, KT, JV and WG. All authors read and approved the final manuscript.

Acknowledgements

This protocol and the forthcoming review are financed by the Mistra Council for Evidence-based Environmental Management (EviEM). Three anonymous reviewers contributed to significant improvements of the manuscript.

References

1. Smith VH: **Eutrophication of freshwater and coastal marine ecosystems - A global problem.** *Environ Sci Pollut Res* 2003, **10**:126–139.
2. Mitsch WJ, Day JW, Zhang L, Lane RR: **Nitrate-nitrogen retention in wetlands in the Mississippi river basin.** *Ecol Eng* 2005, **24**:267–278.
3. Carleton JN, Grizzard TJ, Godrej AN, Post HE: **Factors affecting the performance of storm water treatment wetlands.** *Water Res* 2001, **35**:1552–1562.
4. Kadlec RH: **Nitrogen farming for pollution control.** *J Environ Sci Health A Tox Hazard Subst Environ Eng* 2005, **40**:1307–1330.
5. Vymazal J: **Removal of nutrients in various types of constructed wetlands.** *Sci Total Environ* 2007, **380**:48–65.
6. Vymazal J, Kroepfelova L: **Removal of nitrogen in constructed wetlands with horizontal sub-surface flow: a review.** *Wetlands* 2009, **29**:1114–1124.
7. Hoffmann CC, Kjaergaard C, Uusi-Kamppa J, Hansen HCB, Kronvang B: **Phosphorus retention in riparian buffers: review of their efficiency.** *J Environ Qual* 2009, **38**:1942–1955.
8. Braskerud BC, Tonderski KS, Wedding B, Bakke R, Blankenberg AGB, Ulen B, Koskiaho J: **Can constructed wetlands reduce the diffuse phosphorus loads to eutrophic water in cold temperate regions?** *J Environ Qual* 2005, **34**:2145–2155.
9. Mitsch WJ, Zhang L, Stefanik KC, Nahlik AM, Anderson CJ, Bernal B, Hernandez M, Song K: **Creating wetlands: primary succession, water quality changes, and self-design over 15 years.** *Bioscience* 2012, **62**:237–250.
10. Nixon SW, Ammerman JW, Atkinson LP, Berounsky VM, Billen G, Boicourt WC, Boynton WR, Church TM, Ditoro DM, Elmgren R, *et al*: **The fate of nitrogen and phosphorus at the land sea margin of the North Atlantic Ocean.** *Biogeochemistry* 1996, **35**:141–180.
11. Ryther JH, Dunstan WM: **Nitrogen, phosphorus, and eutrophication in coastal marine environment.** *Science* 1971, **171**:1008–1013.
12. Granéli E, Wallström K, Larsson U, Graneli W, Elmgren R: **Nutrient limitation of primary production in the Baltic Sea area.** *Ambio* 1990, **19**:142–151.
13. Conley DJ, Paerl HW, Howarth RW, Boesch DF, Seitzinger SP, Havens KE, Lancelot C, Likens GE: **Controlling eutrophication: nitrogen and phosphorus.** *Science* 2009, **323**:1014–1015.
14. Schindler DW, Hecky RE, Findlay DL, Stainton MP, Parker BR, Paterson MJ, Beaty KG, Lyng M, Kasian SEM: **Eutrophication of lakes cannot be controlled by reducing nitrogen input: results of a 37-year whole-ecosystem experiment.** *Proc Natl Acad Sci USA* 2008, **105**:11254–11258.
15. Boesch D, Hecky R, O'Melia C, Schindler D, Seitzinger S: *Eutrophication of Swedish Seas.* Stockholm: Swedish Environmental Protection Agency, Report 5509; 2006.


16. Danielsson Å, Papush L, Rahm L: **Alterations in nutrient limitations — scenarios of a changing Baltic Sea.** *J Mar Syst* 2008, **73**:263–283.
17. Swedish Meteorological and Hydrological Institute. [<http://vattenwebb.smhi.se/wetlands/>] (in Swedish).
18. Svensson JM, Strand J, Sahlén G, Weisner S: **Utvärdering av våtmarker anlagda inom lokala investeringsprogram och med LBU-stöd avseende närsaltsretention och biologisk mångfald.** In *Rikare mångfald och mindre kväve, Utvärdering av våtmarker skapade med stöd av lokala investeringsprogram och landsbygdsutvecklingsstöd* (in Swedish). Naturvårdsverket, Rapport 5362; 2004.
19. Brandt M, Arheimer B, Gustavsson H, Pers C, Rosberg J, Sundström M, Thorén AK: *Uppföljning av effekten av anlagda våtmarker i jordbrukslandskap. Belastning av kväve och fosfor* (in Swedish). Naturvårdsverket, Rapport 6309; 2009.
20. Weisner S, Thiére G: *Mindre fosfor och kväve från jordbrukslandskapet* (in Swedish). Jordbruksverket, Rapport 2010; 2010:21.
21. Kottek M, Grieser J, Beck C, Rudolf B, Rubel F: **World map of the Koppen-Geiger climate classification updated.** *Meteorol Z* 2006, **15**:259–263.
22. Thiére G, Milenkovski S, Lindgren PE, Sahlén G, Berglund O, Weisner SEB: **Wetland creation in agricultural landscapes: biodiversity benefits on local and regional scales.** *Biol Conserv* 2009, **142**:964–973.
23. Fisher J, Acreman MC: **Wetland nutrient removal: a review of the evidence.** *Hydrol Earth Syst Sci* 2004, **8**:673–685.
24. Mermillod-Blondin F, Lemoine D, Boisson JC, Malet E, Montuelle B: **Relative influences of submersed macrophytes and bioturbating fauna on biogeochemical processes and microbial activities in freshwater sediments.** *Freshw Biol* 2008, **53**:1969–1982.
25. Hahn S, Bauer S, Klaassen M: **Quantification of allochthonous nutrient input into freshwater bodies by herbivorous waterbirds.** *Freshw Biol* 2008, **53**:181–193.
26. Andersen DC, Sartoris JJ, Thullen JS, Reusch PG: **The effects of bird use on nutrient removal in a constructed wastewater-treatment wetland.** *Wetlands* 2003, **23**:423–435.

Mistra EviEM Protocols

SR1 Protocol: **What are the impacts of reindeer/caribou (*Rangifer tarandus L.*) on arctic and alpine vegetation?** A systematic review protocol. (March 2013)

SR2 Protocol: **How effective are created or restored freshwater wetlands for nitrogen and phosphorus removal?** A systematic review protocol. (September 2013)

SR3 Protocol: **What is the influence on water quality in temperate eutrophic lakes of a reduction of planktivorous and benthivorous fish?** A systematic review protocol. (May 2013)



Mistra EviEM conducts systematic reviews of environmental issues identified as important by public agencies and other stakeholders. These provide an overall assessment of the state of scientific knowledge and help to improve the basis for environmental decision-making in Sweden.

EviEM

Mistra Council for Evidence-based Environmental Management

Address:

Royal Swedish Academy of Sciences
Box 50005, SE-104 05 Stockholm, Sweden

Visit/Deliveries:

Lilla Frescativägen 4A, SE-114 18 Stockholm, Sweden

Phone: 08-673 95 00 | Email: info@eviem.se | www.eviem.se